

Biphasic predators provide biomass subsidies in small freshwater habitats: A case study of spring and cave pools

Benedetta Barzagli | Gentile Francesco Ficetola | Roberta Pennati | Raoul Manenti 

Department of Environmental Science and Policy, Università degli Studi di Milano, Milan, Italy

Correspondence

Raoul Manenti, Department of Environmental Science and Policy, Università degli Studi di Milano, Milano, Italy.
Email: raoulmanenti@gmail.com

Summary

1. There is increasing attention on how biomass exchanges determine linkages between ecosystems. However, there is limited information on the factors determining the direction and overall magnitude of energy flow, especially in systems where primary producers are limited.
2. In this study we compared two aquatic ecosystems (cave and spring pools) using the biomass exchanges driven by a biphasic predator, the fire salamander (*Salamandra salamandra*).
3. Between 2013 and 2014, we monitored 21 fire salamander breeding sites (12 cave pools and 9 spring pools) and we quantified the larval biomass input–export. The balance between input and export was related to different abiotic and biotic variables.
4. *Salamandra salamandra* larvae constituted a major component of the animal biomass in both cave and spring pools, and were the most abundant top predator. Light was the most important parameter in constraining predator biomass; biomass export was lowest in the sites that received less light. In cave pools, the biomass balance was negative, while it was positive in spring pools.
5. Our study demonstrates that cave pools are a system with much lower productivity than springs, and their functioning strongly depends on the input of external resources. Predator occurrence may constitute a major trophic subsidy in poorly productive environments. Gradients of abiotic variables such as light incidence can constrain population size of predators and determine a large variation in their biomass balance.

KEYWORDS

biomass, biospeleology, energy flow, *Salamandra*, subsidy

1 | INTRODUCTION

Biomass production and exchange are key properties in determining ecosystem functioning (Wallace, Eggert, Meyer, & Webster, 2015). General theoretical models affirm that all the trophic levels of an ecosystem have a positive response to productivity gradients (Chase, 2003). Resource supply is a fundamental constraint on the biomass and production at all trophic levels. Unravelling the processes that determine the amount of available resources in ecosystems is pivotal

in understanding their functioning, and also allows assessment of the role of environmental conditions that affect the outcome of complex biotic interactions (Kiffney, Richardson, & Bull, 2003; Knapp, 2005; Kupisch, Moenickes, Schlieff, Frassl, & Richter, 2012). One aspect that is gaining increasing attention in functional ecology is how the exchanges from surrounding ecosystems may affect consumer and predator biomass and abundance in certain environments (Iskali & Zhang, 2015; Venarsky, Huntsman, Huryn, Benstead, & Kuhajda, 2014). Exchanges of detritus, trophic resources, prey and predators

between nearby ecosystems play important roles in the functioning of ecosystems (Kreps, Larson, & Lodge, 2016; Regester, Lips, & Whiles, 2006).

Spatial and functional linkages between two ecosystems may occur because of abiotic processes that spread or displace organic matter, and through the active or passive moving of organisms between different habitats (Kraus, Pletcher, & Vonesh, 2011). Many taxa exploit multiple ecosystems during their life cycle. For instance, several lucifugous organisms exploit both underground and outdoor habitats during their life cycles. The movements of these animals between these two environments likely have strong effects on the fluxes of resources between these adjacent ecosystems (Fenolio, Graening, Collier, & Stout, 2006; Fenolio, Graening, & Stout, 2005; Iskali & Zhang, 2015; Lunghi, Manenti, & Ficetola, 2014). Outside caves, species with biphasic life cycles have a comparable role. For example, many insects and amphibians lay eggs and larvae in water. After metamorphosis they show fully terrestrial adult stages (Earl & Semlitsch, 2012; Eitam, Blaustein, & Mangel, 2002), and the flow of individuals can determine a relevant flow of biomass between terrestrial and aquatic ecosystems. Such a flow can be positive (i.e. with biomass input into aquatic ecosystems higher than biomass export to terrestrial environments), or negative, with input lower than export. Despite the importance of biomass flow for small and nutrient-poor ecosystems, there is limited information on the factors determining the direction and overall magnitude of such flows (Reinhardt, Steinfartz, Paetzold, & Weitere, 2013; Schneider, Christman, & Fagan, 2011).

Ecosystems in which light availability is a constraining factor may provide new insights into some of these interactions. Recent studies show that algal production supports most of the animal production in small freshwater ecosystems (Brett et al., 2017). Studies performed in heavily shaded headwater ecosystems have underscored that primary production and consumer biomass have a positive relationship with light flux (Hill, Ryon, & Schilling, 1995; Kiffney et al., 2003). Therefore, variation in light has been proposed as an important factor affecting the strength of food limitation. For example, even a limited decrease in light availability can strongly reduce the abundance of shredders, the biomass of fungi and the rate of litter breakdown in streams (Laguer et al., 2011). However, the relationships between light intensity and the abundance of different taxa may be complex, for instance, because light flux may interact with the abundance of top predators, thus constraining the abundance at lower trophic levels (Kiffney, 2008).

Caves represent a system where local production and allochthonous inputs are easily discernible. Lack of light determines heterotrophic food webs completely dependent on allochthonous inputs (Romero, 2009; Schneider et al., 2011). When allochthonous detritus subsidises a habitat, the type of response displayed by the organisms depends on the trophic level that receives the input, on the resource features (Mammola & Isaia, 2016) and also on the abiotic features of the habitat itself (Schneider, Kay, & Fagan, 2010; Schneider et al., 2011).

Although cave habitats are often reported as ideal donor-controlled habitats (Polis et al. 1997), few studies have analysed the

effects of subsidies on cave communities (Schneider et al., 2011) and no studies have analysed the possible subsidy effect caused by the predator trophic level. The types of trophic inputs entering cave environments are highly variable and include dead leaves and wood (Pipan, Navodnik, Janzekovic, & Novak, 2008; Romero, 2009), and the faecal material deposited by bats and insects (Fagan, Lutscher, & Schneider, 2007). Moreover, organisms entering caves may themselves be an input of energy for the system (Lunghi et al., 2014; Manenti, 2014).

Salamanders have a particularly important role in small aquatic systems (Davic & Welsh, 2004). The European fire salamander (*Salamandra salamandra*: Salamandridae) is a predator with a terrestrial adult phase and an aquatic larval phase (Krause & Caspers, 2015; Manenti, Ficetola & De Bernardi, 2009; Steinfartz, Stemshorn, Kuesters, & Tautz, 2006). These salamanders generally breed in streams (Denoël & Winandy, 2014), and may have significant effects on subsidy exchanges between pools and surrounding woods (Reinhardt et al., 2013). The species is usually ovoviviparous, and adult salamanders feed exclusively in the terrestrial habitat, thus the biomass of laid larvae does not depend on the resources of the aquatic ecosystem where they live (Reinhardt et al., 2013). Therefore, birth and metamorphosis of larvae determine consistent bidirectional fluxes of biomass between aquatic breeding sites and the terrestrial environment (Reinhardt et al., 2013). The fire salamander has highly plastic breeding strategies, and in karst areas it also actively uses cave pools as its breeding habitat (Manenti, Ficetola, Bianchi, & De Bernardi, 2009; Manenti, Ficetola, Marieni, & De Bernardi, 2011). Salamanders are able to successfully develop in cave pools, but the underground environment strongly affects larval development, which in caves requires more time to reach metamorphosis than in streams (Limongi, Ficetola, Romeo, & Manenti, 2015; Manenti et al., 2011). The cave waterbodies are predator free, therefore fire salamander larvae are the top predators in this system. Similar to what occurs in streams, larvae deposited in cave pools represent an intake of biomass from the outdoor terrestrial environments for cave pools, while metamorphosing individuals export biomass from the underground aquatic environment to the surrounding terrestrial one.

The aim of this study was to assess the factors determining the balance of biomass in cave and spring pools. In particular, we examined (1) how light intensity, together with other environmental features, drives biomass export and (2) if top predators constitute a trophic subsidy in these ecosystems. These aspects provide new insights into the role played by predators in the energy flows of small freshwater systems.

2 | MATERIALS AND METHODS

From March to August 2013, and from March to August 2014, we monitored 21 fire salamander breeding sites (12 cave pools and 9 spring pools). Sites were relatively isolated spring or cave pools (Figure S1). Both cave and spring pools were headwater, therefore compensatory drift (Krause & Caspers, 2015; Reinhardt et al., 2013) of

fire salamander larvae was impossible. In both habitats pools were fed by water from the aquifer and showed limited flow, being distant from eventual downstream running freshwater habitats. All the sites were located in Lombardy (NW Italy) in the Pre-alpine region (Figure 1). Each site was surveyed weekly, for a total of 24 surveys per site during the same period; each site was monitored during one single year; 2 years were necessary to follow all the sites. In 2013 we surveyed six cave and five spring pools, while in 2014 we surveyed six cave and four spring pools.

In each site, fire salamander larvae were caught using a small dip net (mesh size: 1 mm). Catching was performed until no larvae were visible in the site, and was continued for at least 10 min after capturing the last larva. Larvae were placed in a small aquarium (20 × 10 cm, depth 5 cm), and the capture session was repeated 30 min later. For each larva, we recorded total length (accuracy: 1 mm) and weight (accuracy: 0.01 g). Furthermore, following the recommendation of Eitam and Blaustein (2002), the larvae were photographed laterally with a Casio Exilim Ex H30 camera to allow individual identification on the basis of tail colour pattern. The abundance of salamanders for each site and sampling session was estimated on the basis of removal sampling. Removal sampling is an approach to population size estimation that requires the systematic capture and removal of individuals. Population size can be estimated on the basis of the decline in catch size during the sequential sessions of capture and catch efforts. Population size was estimated using the homogeneous capture probability model proposed by Chao and Chang (1999), which provides reliable estimates of the number of individuals present.

The quantification of biomass balance was obtained by comparing the biomass of larvae deposited in the spring to the biomass of metamorphosing larvae during the period studied, which represents the biomass export from each site through salamanders (Reinhardt

et al., 2013). On the whole, we collected and weighed more than 1,540 larvae. To quantify the larval biomass input, we first calculated the maximum number of newborn larvae laid in each site on the basis of the removal method and then multiplied the average body weight of newborns at a given site by the estimated number of newborn larvae (Reinhardt et al., 2013). Fire salamander larvae stop feeding at late development stages (Reinhardt et al., 2013). Therefore, to estimate biomass export we considered the weight of all the fire salamander larvae (distinguished individually on the basis of their lateral colour pattern) that reached the 3B pre-metamorphosis developmental stage (Jusczyk & Zakrzewski, 1981) at a given site (Figure S2). For estimating biomass export, we considered the total weight of individuals collected through removal samplings at the pre-metamorphosis stage; at the 3B stages salamander larvae have already established their aposematic coloration and their detectability is very high, thus limiting the risk of overlooking individuals. We considered as exported biomass the weight recorded for 3B larvae that were not observed in successive surveys. The sum of these weights was considered as the exported biomass at the fire salamander larvae level.

We measured both abiotic and biotic variables potentially related to the balance of biomass export at fire salamander level. At each survey and for each pool, we recorded surface area on the basis of pool maximum width and length, we measured the maximum water level and recorded water temperature measured in the middle of the pool at 2 cm from the substrate during the same daytime range (10–12 a.m.). We also recorded the maximum illuminance incident on the pool surface using a CEM DT8820 multiparameter (range 0.01–200,000 lux). As biotic parameters we considered measures of predation pressure (biomass of predators), intraspecific competition (abundance of salamander larvae) and an estimate of prey availability (biomass of potential prey).




FIGURE 1 Study area (map shows Italian political borders). Triangles show locations of cave pools; circles show locations of spring pools. Some of the symbols are superimposed owing to geographic proximity

Macroinvertebrate biomass was measured using pipe sampling (Dodd, 2010). Samples were collected by thrusting a 0.3 m² circular pipe sampler through the water column and about 5 cm into the sediment. Small nets (mesh size: 1 mm) were used to remove all animals from the water and the first centimetre of the sediment (Dodd, 2010). Net sweeps were collected until at least 10 consecutive sweeps were empty; for each site we repeated pipe sampling twice. Once captured, we weighed macroinvertebrates on a digital PESOLA balance (precision 0.01 g), by distinguishing between potential prey and predators of fire salamander larvae, on the basis of taxonomy and dimensions. As predators, we considered freshwater crayfish longer than 35 mm and dragonfly larvae longer than 20 mm (Bo, Fenoglio, Lopez-Rodriguez, & De Figueroa, 2011; Ghia, Fea, Spairani, Bernini, & Nardi, 2009; Manenti, Siesa, & Ficetola, 2013). Smaller macroinvertebrates were considered as prey; we did not record crayfish <35 mm. We then estimated prey and predator wet biomass for each site (g/m²).

We used linear models (LMs) to assess the factors determining the balance of biomass export, which was considered as a dependent variable: as independent variables we considered average maximum illuminance recorded during the whole period, site type (cave/spring), average water temperature, fire salamander average density, predator average biomass and prey average biomass. Salamander density and export biomass were not correlated ($r = .20$; $t = 0.86$, $p = .39$), thus suggesting that the two variables are essentially independent. If needed, independent variables were transformed using a logarithm (illuminance, larvae density, prey and predator biomass) to reduce skewness and improve normality. We built models representing all the possible combinations of independent variables and ranked them on the basis of corrected Akaike's Information Criterion (AICc) (Rolls, 2011). As AICc may select overly complex models, we considered a complex model only if it showed AICc less than the AICc of all of its simpler nested models (Richards, Whittingham, & Stephens, 2011). Models expressing the highest proportion of variation using the smallest number of predictors have the smallest AICc values and are considered the "best models" (Lukacs et al., 2007). We also assessed significance of variables composing the best model using a likelihood ratio test (Bolker et al., 2008). We performed all the analyses in R 3.31 environment using the, nlme, car and MuMIn packages.

3 | RESULTS

We recorded fire salamander larvae metamorphosis events in all the cave and spring pools. Cave pools were generally dark, but light partially hit some of the pools (average \pm SE illuminance of the most luminous part of cave pools: 54.3 ± 34 lux; range 0.00–400 lux), particularly in sites where the pool used for larvae deposition was close to the cave entrance. Spring pools were much more luminous, ranging from 10,100 to 45,300 lux (average \pm SE: $17,171 \pm 4,073$ lux). During the study period, cave pools showed a noticeable stability in terms of constancy of the surface area,

without appreciable variations. Instead, spring pools varied much more, with a maximum surface reduction of 95% (average \pm SE total variation in surface in epigeal sites: $13.34\% \pm 4\%$); a Leven's homogeneity test, performed on the surface of pools between two consecutive surveys, confirmed that the surface of spring pools undergoes much stronger variations than does the surface of cave pools ($F_{1,61} = 10.28$, $p < .01$).

No predators were detected in caves, except for one site in which we recorded two juveniles of the native crayfish *Austropotamobius pallipes* (Astacidae), while in spring pools predators occurred in almost all the sites ($n = 7$) and were mostly composed of *Cordulegaster* (Cordulegasteridae) and *Aeshna* (Aeshnidae) dragonfly larvae (maximum density 2.09 individuals/m²; mean density \pm ES = 1.39 ± 0.37 individuals/m²).

In March, the average biomass (\pm SE) of newborn fire salamander larvae was 4.91 ± 1.93 g/m in caves and 19.9 ± 6.10 g/m in spring pools. The difference in biomass import per square metre was weakly significant ($t_{1,9} = 2.34$, $p = .04$). In cave pools the average biomass export was 40% of the import, while in spring pools it was 73% of the import. Maximum salamander larvae biomass in cave pools was 53.3 g/m, representing on average 97.3% of the total aquatic biomass of the pool. Conversely, in spring pools the maximum biomass of the larvae was 17.3 g/m, with salamander larvae accounting for 27.6% of the total biomass (Figure 2).

The biomass balance was negative for 75% of cave pools, while all the spring pools had a positive balance (Table 1). According to the best-AICc model, light intensity was the most likely determinant of biomass balance, with a strong, negative relationship between biomass balance and light ($F_{1,17} = 20.56$, $p < .001$). Although illuminance and site type were highly correlated ($r = .83$, $t_{1,18} = 6.4$, $p < .0001$), site type (cave vs. spring) was not included among the best AIC models, and the importance of light intensity was much higher than that of site type. In the best selected model, larvae density was also included, but the relationship was weak and not significant ($F_{1,17} = 1.33$, $p = .26$). The importance of all the remaining predictors, including predator and prey biomass, was extremely low (Tables 2 and 3).

4 | DISCUSSION

Fire salamander larvae constitute a major component of the animal biomass in both cave and spring pools, and are the most abundant top predators for them. Therefore, identifying the factors determining their biomass balance is pivotal to understanding the functioning of the small freshwater ecosystems in which they live. Abiotic environmental features were more important than biotic parameters in determining salamander biomass balance, as biomass production was generally highest in the sites that received more light and the most productive ones for the larvae. These results underscore that light flux is a main constraint on predator biomass; a similar pattern, which applies only to lower trophic levels, has been reported for shaded headwater streams in which

FIGURE 2 Box-whisker plots of biomass levels recorded in spring and cave pools. The bold bar represents the median; upper and lower box bars represent the upper and lower quartile, respectively; whiskers represent the maximum values. (a) Comparison between the mean biomass of salamander larvae and their predators and prey in epigeal sites; (b) Comparison between the mean biomass of salamander larvae and their predators and prey in cave pools; (c) Comparison of salamander larvae biomass import between spring and cave pools; (d) Comparison of salamander larvae biomass export between spring and cave pools




TABLE 1 Environmental features of the spring and cave pools surveyed

ID site	Pool type	Import (g)	Export (g)	Biomass balance (g)	Average temperature (°C)	Average maximum illuminance recorded (lux)	Average larvae density (g/m ²)	Average prey biomass (g)	Average predator biomass (g/m ²)
G01	Cave	3.58	0	-3.58	13.4	451	2.41	0.08	0
G02	Cave	24.47	24.5	0.03	11.7	26.6	25.1	0.11	0
G03	Cave	16.74	0	-16.74	11.2	0.89	2.5	0	0
G07	Cave	0.99	1.35	0.36	12.8	9.6	5.63	0.007	0
G09	Cave	5.34	5.96	0.62	11	3.96	3.9	0.18	0
G11	Cave	12.12	0	-12.12	12.1	5.8	1.76	0	1.19
G16	Cave	11.85	4.23	-7.62	14.3	53	7.86	0.13	0
G17A	Cave	10.57	3.52	-7.05	12.1	16.56	1.12	0.1	0
G17B	Cave	3.08	0	-3.08	11.6	17	0.3	0.05	0
G17C	Cave	1.1	0	-1.1	11.7	17	2.35	0.032	0
G19	Cave	5.47	0	-5.47	10.16	0	0	1.258	0
G22	Cave	8.78	2.28	-6.5	12.2	24.3	1.6	0.085	0
Pozzaa	Spring	56.12	75.46	19.34	11.91	18,300	17.31	0.2	0.28
Pozzab	Spring	18.92	70.08	51.16	15.75	45,300.00	2.41	0.17	0
Pozzac	Spring	86.93	6.15	-80.78	11.26	10,100	14.39	0.1	0
Pozzad	Spring	91.78	8.66	-83.12	15.55	5300	46.74	3.26	1.89
Pozza1	Spring	1.16	3.2	2.04	17.22	16,125	22.9	0.12	0.13
Pozza2	Spring	9.43	11.18	1.75	17.31	14,875	26.09	0.4	1.76
Pozza3	Spring	14.28	16.62	2.34	15.2	9,875	76.24	0.24	0
Pozza4	Spring	23.17	27.03	3.86	16.67	18,666	34.17	3.19	2.4
Pozza5	Spring	3.53	7.11	3.58	18.36	16,000	29.5	0.06	0.9

primary producer and consumer biomasses are constrained by light flux with consequences for nutrient supply and predator abundance (Kiffney, 2008). Primary producers, such as periphyton, photosynthetic bacteria occurring in biofilms, and other photosynthetic organisms, are strongly limited in shady environments (Chase, 2003). Several studies in shaded headwaters have shown that light

flux increases primary production and consumer biomass (Hill et al., 1995; Kiffney, 2008), even though high light flux over a certain level can limit primary production through photoinhibition (Hill et al., 1995; Kiffney et al., 2003; Murray, Tenhunen, & Nowak, 1993). Although almost all the study sites received some light, the comparison between spring and cave pools with great

TABLE 2 Best models with AICc weight > 0.01 illustrating the relationship between biomass export and the environmental variables of the cave and spring pools

Variables in the best models	R ²	AICc	Δ	Weight
Illuminance, larvae density	0.56	155.2	0.00	0.59
Illuminance	0.41	157.8	2.53	0.17
Larvae density, cave	0.49	158.0	2.82	0.14
Cave	0.36	159.5	4.24	0.07
Average temperature	0.29	161.4	6.22	0.03
Null model	0.00	165.6	10.38	0.00
Larvae density	0.03	167.7	12.47	0.00

TABLE 3 Model-averaged parameters of environmental variables influencing the biomass balance. *w*: cumulative AICc weight of the variable; *B*: averaged regression coefficient, with ±standard error

Variable	<i>w</i>	<i>B</i>	±SE
Illuminance	0.76	-7.25	1.66
Larvae density	0.73	13.75	0.19
Cave	0.21	17.01	0.00
Average temperature	0.03	-5.20	0.00

differences in light, clearly reveals the dominating effects of this parameter.

In cave pools, the biomass balance was generally negative, with a pattern opposite to that observed in springs. Our study demonstrates that cave aquatic environments, even when close to cave entrances, are clearly less subsidising ecosystems than outdoor springs, even though some of them occur within heavily shaded forests. This may be linked to different factors. First, in headwater streams and springs, the occurrence of periphyton biofilms and other primary producers increases the availability of nutrients for the food web together with the detritus and organic matter deriving from the surrounding terrestrial habitat (Brett et al., 2017). Recent studies demonstrate that the biomass of benthic algae (such as diatoms), although relatively minor compared to detritus, represents important high-quality food sources for some of the invertebrates that are fire salamander prey (Crenier et al., 2017).

Second, epigeal sites may be more easily accessible to biphasic insects, the larvae of which often are detritivore or primary consumers (Koperski, 2011) and constitute the usual prey of the fire salamander larvae (Costa, Salvidio, Romano, & Baroni, 2015). The adult females of salamanders feed in terrestrial habitats such as the forests surrounding breeding sites. Our results indicate that when salamanders enter caves to deposit their larvae, they deliver a major biomass subsidy to these habitats, and such a biomass input is much higher than the export that will return to the outdoor environment. This represents an interesting case of a direct subsidy from the top predator level. In this extreme case, top predators, although exerting strong predation pressures on consumers at lower trophic levels, constitute the major biomass input of the system, and thus have a

major role in sustaining the whole food web. The importance of salamander larvae as an aquatic food-web subsidy has been observed in other systems, such as in low-productivity temporary ponds in forests, where the amount of larval biomass that enters the ponds is generally higher than the export through metamorphosed individuals (Reinhardt et al., 2013).

The behaviour of the fire salamander actively entering caves to breed is a recently described phenomenon (Manenti, Ficetola, Bianchi et al., 2009), which may be favoured by the fact that cave habitats in temperate regions are generally predator-free environments (Fiser, Blejec, & Trontelj, 2012; Galassi, Stoch, Fiasca, Di Lorenzo, & Gattone, 2009; Schneider et al., 2011) and provide stable environmental conditions. However, cave environments also pose multiple limitations, and food scarcity severely limits growth rates (Limongi et al., 2015). As a consequence, a particularly high reproductive effort is needed to exploit underground fresh waters successfully. Breeding fire salamander females provide a substantial contribution to biomass inputs in both spring and cave pools and can subsidise heterotrophic systems in which cannibalism is likely to play a major role. Both in cave and spring pools, it is frequent to find different cohorts of larvae showing strong asymmetries in terms of both size and developmental stage. The smallest salamander larvae, belonging to the youngest cohorts, may represent an important trophic supply for the larger conspecifics (Limongi et al., 2015; Romeo, Giovine, Ficetola, & Manenti, 2015). In the case of the study pools, most of the prey-predator energy flow likely occurs at the same trophic level, i.e. among top predators. The evolutionary adaptations allowing the exploitation of different environments by a single species may thus be driven by female choice of breeding sites, and can result in a change in the direction of energy fluxes between aquatic and terrestrial environments, as also reported for temporary pools (Reinhardt et al., 2013; Steinfartz, Weitere, & Tautz, 2007).

Overall, our study highlights two key points. First, the exploitation of a habitat by the temporary life stages of animals at the predator level does not necessarily translate into a depletion of the general biomass of an ecosystem; on the contrary, in poorly productive environments, it may constitute a trophic subsidy. Second, the biomass balance of predators may vary as a result of abiotic gradients, such as light intensity, that constrain population size.

ACKNOWLEDGMENTS

We are grateful to G. Romeo, L. Limongi and M. Toma for help during field surveys. The suggestion of T. Mehner and two anonymous reviewers significantly improved a previous version of the manuscript.

REFERENCES

- Bo, T., Fenoglio, S., Lopez-Rodriguez, M. J., & De Figueroa, J. M. T. (2011). Trophic behaviour of the dragonfly *Cordulegaster boltoni* (Insecta: Odonata) in small creeks in NW Italy. *Entomologica Fennica*, 22, 255–261.

- Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., & White, J. S. (2008). Generalized linear mixed models: A practical guide for ecology and evolution. *Trends in Ecology & Evolution*, *24*, 127–135.
- Brett, M. T., Bunn, S. E., Chandra, S., Galloway, A. W. E., Guo, F., Kainz, M. J., ... Wehr, J. D. (2017). How important are terrestrial organic carbon inputs for secondary production in freshwater ecosystems? *Freshwater Biology*, *62*, 833–853.
- Chao, A., & Chang, S. H. (1999). An estimating function approach to the inference of catch-effort models. *Environmental and Ecological Statistics*, *6*, 313–334.
- Chase, J. M. (2003). Strong and weak trophic cascades along a productivity gradient. *Oikos*, *101*, 187–195.
- Costa, A., Salvidio, S., Romano, A., & Baroni, D. (2015). Larval diet of *Salamandra salamandra* (L., 1758): Preliminary results on prey selection and feeding strategy. In G. Doria, R. Poggi, S. Salvidio, & M. Tavano (Eds.), *Atti X congresso Nazionale della Societas Herpetologica Italica (Genova 15–18 Ottobre 2014)* (pp. 33–38). Pescara: Ianieri Edizioni.
- Crenier, C., Arce-Funck, J., Bec, A., Billoir, E., Perrière, F., Leflaive, J., ... Danger, M. (2017). Minor food sources can play a major role in secondary production in detritus-based ecosystems. *Freshwater Biology*, *62*, 1155–1167.
- Davic, R. D., & Welsh, H. H. (2004). On the ecological roles of salamanders. *Annual Review of Ecology Evolution and Systematics*, *35*, 405–434.
- Denoël, M., & Winandy, L. (2014). Fire salamander (*Salamandra salamandra*) in Larzac plateau: Low occurrence, pond-breeding and cohabitation of larvae with paedomorphic palmate newts (*Lissotriton helveticus*). *Acta Herpetologica*, *9*, 43–49.
- Dodd, C. K. J. (2010). *Amphibian ecology and conservation*. Oxford: Oxford University Press.
- Earl, J. E., & Semlitsch, R. D. (2012). Reciprocal subsidies in ponds: Does leaf input increase frog biomass export? *Oecologia*, *170*, 1077–1087.
- Eitam, A., & Blaustein, L. (2002). Noninvasive individual identification of larval *Salamandra* using tailfin spot patterns. *Amphibia-Reptilia*, *23*, 215–219.
- Eitam, A., Blaustein, L., & Mangel, M. (2002). Effects of *Anisops sardea* (Hemiptera: Notonectidae) on oviposition habitat selection by mosquitoes and other dipterans and on community structure in artificial pools. *Hydrobiologia*, *485*, 183–189.
- Fagan, W. F., Lutscher, F., & Schneider, K. (2007). Population and community consequences of spatial subsidies derived from central-place foraging. *The American Naturalist*, *170*, 902–915.
- Fenolio, D. B., Graening, G. O., Collier, B. A., & Stout, J. F. (2006). Coprophagy in a cave-adapted salamander; the importance of bat guano examined through nutritional and stable isotope analyses. *Proceedings of the Royal Society B: Biological Sciences*, *273*, 439–443.
- Fenolio, D. B., Graening, G. O., & Stout, J. F. (2005). Seasonal movement patterns of pickerel frogs (*Rana palustris*) in an Ozark cave and trophic implications supported by stable isotope evidence. *The Southwestern Naturalist*, *50*, 385–389.
- Fiser, C., Blejcek, A., & Trontelj, P. (2012). Niche-based mechanisms operating within extreme habitats: A case study of subterranean amphipod communities. *Biology Letters*, *8*, 578–581.
- Galassi, D. M. P., Stoch, F., Fiasca, B., Di Lorenzo, T., & Gattone, E. (2009). Groundwater biodiversity patterns in the Lessinian Massif of northern Italy. *Freshwater Biology*, *54*, 830–847.
- Ghia, D., Fea, G., Spairani, M., Bernini, F., & Nardi, P. A. (2009). Movement behaviour and shelter choice of the native crayfish *Austropotamobius pallipes* complex: Survey on a population in a semi-natural pond in Northern Italy. *Marine and Freshwater Behaviour and Physiology*, *42*, 167–185.
- Hill, W. R., Ryon, M. G., & Schilling, E. M. (1995). Light limitation in a stream ecosystem: Responses by primary producers and consumers. *Ecology*, *76*, 1297–1309.
- Iskali, G., & Zhang, Y. X. (2015). Guano subsidy and the invertebrate community in Bracken Cave: The world's largest colony of bats. *Journal of Cave and Karst Studies*, *77*, 28–36.
- Jusczyk, W., & Zakrzewski, M. (1981). External morphology of larval stages of the spotted salamander *Salamandra salamandra* (L.). *Acta Biologica Cracoviensis*, *23*, 127–135.
- Kiffney, P. M. (2008). Response of lotic producer and consumer trophic levels to gradients of resource supply and predation pressure. *Oikos*, *117*, 1428–1440.
- Kiffney, P. M., Richardson, J. S., & Bull, J. P. (2003). Responses of periphyton and insects to experimental manipulation of riparian buffer width along forest streams. *Journal of Applied Ecology*, *40*, 1060–1076.
- Knapp, R. A. (2005). Distribution patterns of lentic-breeding amphibians in relation to ultraviolet radiation exposure in western North America. *Biological Conservation*, *121*, 265–279.
- Koperski, P. (2011). Diversity of freshwater macrobenthos and its use in biological assessment: A critical review of current applications. *Environmental Reviews*, *19*, 16–31.
- Kraus, J. M., Pletcher, L. T., & Vonesh, J. R. (2011). Variation in active and passive resource inputs to experimental pools: Mechanisms and possible consequences for food webs. *Freshwater Biology*, *56*, 491–502.
- Krause, E. T., & Caspers, B. A. (2015). The influence of a water current on the larval deposition pattern of females of a diverging fire salamander population (*Salamandra salamandra*). *Salamandra: German Journal of Herpetology*, *51*, 156–160.
- Kreps, T. A., Larson, E. R., & Lodge, D. M. (2016). Do invasive rusty crayfish (*Orconectes rusticus*) decouple littoral and pelagic energy flows in lake food webs? *Freshwater Science*, *35*, 103–113.
- Kupisch, M., Moenickes, S., Schlieff, J., Frassl, M., & Richter, O. (2012). Temperature-dependent consumer-resource dynamics: A coupled structured model for *Gammarus pulex* (L.) and leaf litter. *Ecological Modelling*, *247*, 157–167.
- Lagrange, C., Kominoski, J. S., Danger, M., Baudoin, J. M., Lamothe, S., Lambriquet, D., & Lecerf, A. (2011). Experimental shading alters leaf litter breakdown in streams of contrasting riparian canopy cover. *Freshwater Biology*, *56*, 2059–2069.
- Limongi, L., Ficetola, G. F., Romeo, G., & Manenti, R. (2015). Environmental factors determining growth of salamander larvae: A field study. *Current Zoology*, *61*, 421–427.
- Lukacs, P. M., Thompson, W. L., Kendall, W. L., Gould, W. R., Doherty, P. F., Burnham, K. P., & Anderson, D. R. (2007). Concerns regarding a call for pluralism of information theory and hypothesis testing. *Journal of Applied Ecology*, *44*, 456–460.
- Lunghi, E., Manenti, R., & Ficetola, G. F. (2014). Do cave features affect underground habitat exploitation by non-troglobite species? *Acta Oecologica*, *55*, 29–35.
- Mammola, S., & Isaia, M. (2016). Step back! Niche dynamics in cave-dwelling predators. *Acta Oecologica*, *75*, 35–42.
- Manenti, R. (2014). Role of cave features for aquatic troglomorphic fauna occurrence: Effects on “accidentals” and troglomorphic organisms distribution. *Acta Zoologica Academiae Scientiarum Hungaricae*, *60*, 257–270.
- Manenti, R., Ficetola, G. F., Bianchi, B., & De Bernardi, F. (2009). Habitat features and distribution of *Salamandra salamandra* in underground springs. *Acta Herpetologica*, *4*, 143–151.
- Manenti, R., Ficetola, G. F., & De Bernardi, F. (2009). Water, stream morphology and landscape: Complex habitat determinants for the fire salamander *Salamandra salamandra*. *Amphibia-Reptilia*, *30*, 7–15.
- Manenti, R., Ficetola, G. F., Marieni, A., & De Bernardi, F. (2011). Caves as breeding sites for *Salamandra salamandra*: Habitat selection, larval development and conservation issues. *North-Western Journal of Zoology*, *7*, 304–309.

- Manenti, R., Siesa, M. E., & Ficetola, G. F. (2013). Odonata occurrence in caves: Active or accidentals? A new case study. *Journal of Cave and Karst Studies*, 75, 205–209.
- Murray, K. J., Tenhunen, J. D., & Nowak, R. S. (1993). Photoinhibition as a control on photosynthesis and production of *Sphagnum* mosses. *Oecologia*, 96, 200–207.
- Pipan, T., Navodnik, V., Janzekovic, F., & Novak, T. (2008). Studies of the fauna of percolation water of Huda luknja, a cave in isolated karst in northeast Slovenia. *Acta Carsologica*, 37, 141–151.
- Polis, G. A., Anderson, W. B., & Holt, R. D. (1997). Toward an integration of landscape and food web ecology: The dynamics of spatially subsidized food webs. *Annual Review of Ecology and Systematics*, 28, 289–316.
- Regester, K. J., Lips, K. R., & Whiles, M. R. (2006). Energy flow and subsidies associated with the complex life cycle of ambystomatid salamanders in ponds and adjacent forest in southern Illinois. *Oecologia*, 147, 303–314.
- Reinhardt, T., Steinfartz, S., Paetzold, A., & Weitere, M. (2013). Linking the evolution of habitat choice to ecosystem functioning: Direct and indirect effects of pond-reproducing fire salamanders on aquatic-terrestrial subsidies. *Oecologia*, 173, 281–291.
- Richards, S. A., Whittingham, M. J., & Stephens, P. A. (2011). Model selection and model averaging in behavioural ecology: The utility of the IT-AIC framework. *Behavioral Ecology and Sociobiology*, 65, 77–89.
- Rolls, R. J. (2011). The role of life-history and location of barriers to migration in the spatial distribution and conservation of fish assemblages in a coastal river system. *Biological Conservation*, 144, 339–349.
- Romeo, G., Giovine, G., Ficetola, G. F., & Manenti, R. (2015). Development of the fire salamander larvae at the altitudinal limit in Lombardy (north-western Italy): Effect of two cohorts occurrence on intraspecific aggression. *North-Western Journal of Zoology*, 11, 234–240.
- Romero, A. (2009). *Cave biology*. New York: Cambridge University Press.
- Schneider, K., Christman, M. C., & Fagan, W. F. (2011). The influence of resource subsidies on cave invertebrates: Results from an ecosystem-level manipulation experiment. *Ecology*, 92, 765–776.
- Schneider, K., Kay, A. D., & Fagan, W. F. (2010). Adaptation to a limiting environment: The phosphorus content of terrestrial cave arthropods. *Ecological Research*, 25, 565–577.
- Steinfartz, S., Stemshorn, K., Kuesters, D., & Tautz, D. (2006). Patterns of multiple paternity within and between annual reproduction cycles of the fire salamander (*Salamandra salamandra*) under natural conditions. *Journal of Zoology*, 268, 1–8.
- Steinfartz, S., Weitere, M., & Tautz, D. (2007). Tracing the first step to speciation: Ecological and genetic differentiation of a salamander population in a small forest. *Molecular Ecology*, 16, 4550–4561.
- Venarsky, M. P., Huntsman, B. M., Huryn, A. D., Benstead, J. P., & Kuhauda, B. R. (2014). Quantitative food web analysis supports the energy-limitation hypothesis in cave stream ecosystems. *Oecologia*, 176, 859–869.
- Wallace, J. B., Eggert, S. L., Meyer, J. L., & Webster, J. R. (2015). Stream invertebrate productivity linked to forest subsidies: 37 stream-years of reference and experimental data. *Ecology*, 96, 1213–1228.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Barzaghi B, Ficetola GF, Pennati R, Manenti R. Biphasic predators provide biomass subsidies in small freshwater habitats: A case study of spring and cave pools. *Freshwater Biol.* 2017;00:1–8. <https://doi.org/10.1111/fwb.12975>